

Tryptic Soy Agar + Lactamator 500 IU + Neutralizing (Irradiated)

General purpose medium for environmental monitoring with inactivation of β -lactam antibiotics and disinfectants..

TYPICAL FORMULA	(g/l)
Casein Peptone	15.0
Soy Peptone	5.0
Sodium Chloride	5.0
Agar	15.0
Histidine	1.0
Lecithin	0.7
Polysorbate 80	5.0
Sodium Thiosulfate	0.5
Lactamator	500 IU
Final pH 7.3 \pm 0.2	

DESCRIPTION

Tryptic Soy Agar + Lactamator 500 IU + Neutralizing (Irradiated) is a general purpose medium in RODAC (Replicate Organisms Detection and Counting) used for environmental monitoring with inactivation of β -lactam antibiotics (penicillins, cephalosporins and carbapenems) and disinfectants.

These gamma-irradiated, triple-bagged plates are particularly suitable for use in restricted areas like isolators and clean rooms.

PRINCIPLE

Casein peptone and soy peptone provide amino acids, nitrogen, carbon, minerals, vitamins and other nutrients which support the growth of microorganism. Sodium chloride maintains the osmotic balance of the medium. Agar is the solidifying agent. Histidine inactivates aldehydes. Lecithin neutralizes quaternary ammonium compounds. Polysorbate 80 (Tween 80) is effective against phenolic compounds and mercurial derivatives. Sodium thiosulfate neutralizes halogen compounds. Lactamator is a mixture of Penicillinase and Cephalosporinase, designed for the inactivation of a wide range of beta-lactam antibiotics.

- 1 International Unit (IU) is defined as the amount of enzyme needed to hydrolyze 1 μ mole of Penicillin G (Penicillinase) or 1 μ mole of Cephalosporin C (Cephalosporinase) per minute at pH 7.0 at 25°C.

TECHNIQUE

RODAC plates are recommended for use in air sampling equipment as well as for surface sampling. Selected surfaces are sampled by firmly pressing the agar medium against the test area. Plates are intended for personnel hygiene monitoring (clothing, gloves or hands) as well.

For detection of bacteria incubate the plates at 30-35°C for 18-72 hours.

For detection of yeasts and moulds incubate at 20-25°C for 2-7 days.

INTERPRETATION OF RESULTS

Observe daily for the formation of colonies.

STORAGE AND TRANSPORT CONDITIONS

2-8°C away from light, until the expiry date on the label. However, our stability studies have shown that the transport at 18-25°C for 4 days, or at 35-39°C for 48 hours, does not alter in any way the performance of the product. Eliminate if signs of deterioration or contamination are evident.

WARNING AND PRECAUTIONS

The product does not contain hazardous substances in concentrations exceeding the limits set by current legislation and therefore is not classified as dangerous. It is nevertheless recommended to consult the safety data sheet for its correct use. The product must be used by properly trained operators only.

DISPOSAL OF WASTE

Disposal of waste must be carried out according to the national and local regulations in force.

REFERENCES

1. Swanson, K.J., F.F. Busta, E.H. Peterson, and M.G. Johnson (1992). Colony Count Methods, p. 75-95.
2. USP 33 – NF 28 (2011) <62> Microbiological examination of non-sterile products: Test for specified microorganisms. Harmonised Method. The United States Pharmacopeial Convention, Rockville, MD. USA.
3. USP 33 – NF 28 (2011) <1116> Microbiological evaluation of clean rooms. The United States Pharmacopeial Convention, Rockville, MD. USA.
4. European Pharmacopeia 7.0 (2011) 7th ed. Chapters 2.6.13. Microbiological examination of non-sterile products: Test for specified microorganisms. Harmonised Method. Council of Europe Strasbourg, France.



LIOFILCHEM® S.r.l.

Via Scozia, Zona Ind.le - 64026, Roseto degli Abruzzi (TE) - ITALY

Tel +39 0858930745 Fax +39 0858930330 Website: www.liofilchem.net E-mail: liofilchem@liofilchem.net

PRODUCT SPECIFICATIONS

NAME

Tryptic Soy Agar + Lactamator 500 IU + Neutralizing (Irradiated)

PRESENTATION

Ready-to-use plates RODAC plates (55 mm) containing 17 ± 1 ml of medium

STORAGE

2-8°C

PACKAGING

Ref.	Content	Packaging
15341S	20 plates	<ul style="list-style-type: none"> packed one by one in a blister packs of two pieces five blisters wrapped in film thermally welded vacuum bag and cardboard box

pH OF THE MEDIUM

7.3 ± 0.2

USE

Tryptic Soy Agar + Lactamator 500 IU + Neutralizing (Irradiated) is a general purpose medium in RODAC (Replicate Organisms Detection and Counting) used for environmental monitoring with inactivation of β -lactam antibiotics (penicillins, cephalosporins and carbapenems) and disinfectants. Particularly suitable for use in restricted areas

TECHNIQUE

Refer to technical sheet of the product

APPEARANCE OF THE MEDIUM

Clear or slightly opalescent, amber

SHELF LIFE

4 months

QUALITY CONTROL










- Control of general characteristics, label and print
- Sterility control
48 hours and 7 days at $22.5 \pm 2.5^\circ\text{C}$, in aerobiosis
48 hours and 7 days at $32.5 \pm 2.5^\circ\text{C}$, in aerobiosis
- Microbiological control
Inoculum for productivity: 50-100 CFU
Inoculum for evaluation of lactamator activity: 10^4 - 10^5 CFU
Incubation Conditions: 18-24 h at $32.5 \pm 2.5^\circ\text{C}$, 48-72 h at $22.5 \pm 2.5^\circ\text{C}$, in aerobiosis

Microorganism		Growth
<i>Staphylococcus aureus</i>	ATCC® 6538	Good
<i>Escherichia coli</i>	ATCC® 8739	Good
<i>Pseudomonas aeruginosa</i>	ATCC® 9027	Good
<i>Bacillus subtilis</i>	ATCC® 6633	Good
<i>Candida albicans</i> *	ATCC® 10231	Good
<i>Aspergillus brasiliensis</i> *	ATCC® 16404	Good

Control of penicillinase and cephalosporinase activities with disc diffusion method

Microorganism	Specification
<i>Staphylococcus aureus</i> ATCC® 6538	No inhibition by Penicillin G 10 IU, Ampicillin 10 μg , Cefotaxime 30 μg , Cefepime 30 μg and Meropenem 10 μg

TABLE OF SYMBOLS

 Batch code	 Do not reuse	 Manufacturer	 Use by	 Fragile, handle with care
 Catalogue number	 Temperature limitation	 Contains sufficient for <n> tests	 Caution, consult instructions for use	



LIOFILCHEM® S.r.l.

Via Scozia, Zona Ind.le - 64026, Roseto degli Abruzzi (TE) - ITALY
Tel +39 0858930745 Fax +39 0858930330 Website: www.liofilchem.net E-mail: liofilchem@liofilchem.net